



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Numerous amoeboid cells are always present in the uterus among the embryos. These are probably concerned in the nutrition of the embryos, since they may be seen passing through the uterine wall, and the uterus is surrounded by a great quantity of cells filled with yellow granules, probably of food material.

Perhaps the most important developmental point thus far made out is that the peribranchial sacs arise as two well defined ectodermal invaginations on the dorsal side of the embryo.

The results, then, support the conclusions of Kowalevsky, Seeliger, Willey, Hjort and Caullery on this head, and oppose those of Della Valle, van Beneden et Julin, Pizon and Garstang, who hold, in one way and another, that these structures arise from the endoderm.

*Notes on Chelyosoma productum, Stimpson.* F. W. BANCROFT.

An examination of about 20 individuals in the collections of the University of California shows that this western ascidian is quite distinct from its Atlantic and Arctic representative, *C. macleayanum*. Stimpson describes the species as having the disk, which is characteristic of the genus, divided into fourteen plates; but in the individuals examined the number was found to vary from thirteen to twenty. This variability is associated with a muscular system that is quite different from what is found in the other member of the genus. In *C. productum* the systems of short muscles joining adjacent plates are wanting, except around the orifices, and are replaced by a series of fibres extending from near the center of the disk to its periphery and some distance down the sides of the animal. The method of attachment of these muscles is different from that described for any other ascidian. Both ends of every bundle of muscle fibres are firmly attached to little projections of

the inner surface of the test. On these the ectoderm is thrown into deep folds and pockets which greatly increase the surface of contact with the test, so that the muscles which are joined to the inner ends of the ectoderm cells cannot tear them away.

The matrix of the test, like that of some other tunicates, consists of an inner layer of cellulose and an outer one, very distinctly separated from it, which is not cellulose, and which corresponds to the 'yellow layer' of the early authors. In our species it is easily seen that this outer layer is formed from the cellulose matrix by the activity of the mesodermic bladder cells which the latter contains. The first traces of the 'yellow substance' are seen about isolated bladder cells near the outer layer, and all transitions can be traced from this stage until the cell and the yellow substance it has produced are incorporated into the outer layer. The other organs of *Chelyosoma* are of a less exceptional character and clearly show that it is more closely related to *Corella* than to any other genus.

*On the Plan of Development of a Myxinoid.*  
BASHFORD DEAN.

The marked dissimilarity in the development of *Bdellostoma* and *Petromyzon* was noted. In the former a large supply of yolk produces a merocytic condition at a very early stage; The head region of the embryo, appearing first, very much as in *Elasmo-* branches, takes its position near the animal pole; the body region is then laid down, apparently by concretion, in an almost straight line extending in the direction of the yolk pole almost the entire length of the egg. The subsequent growth of the embryo constricts both head and tail from the yolk sac, and in very late stages an embryo of nearly two inches lies coiled within the egg. A preliminary study confirms Professor Price's observations as to the great number of gill slits.